

A history of the exploitation of the Ross Sea, Antarctica

David G. Ainley

H.T. Harvey and Associates, 983 University Avenue, Building D, Los Gatos, CA 95032, USA
(dainley@penguinscience.com)

Received March 2009

ABSTRACT. Recent analyses of anthropogenic impacts on marine systems have shown that the Ross Sea is the least affected stretch of ocean on Earth, although historical effects were not included in those studies. Herein the literature is reviewed in order to quantify the extent of extraction of biological resources from the Ross Sea continental shelf and slope from the start of the 20th century. There was none before that time. An intense extraction of Weddell seals *Leptonychotes weddellii* by the expeditions of the ‘heroic’ period and then by New Zealand to feed sled dogs in the 1950–1980s caused the McMurdo Sound population to decrease permanently. Otherwise no other sealing occurred. Blue whales *Balaenoptera musculus intermedia* were extirpated from waters of the shelf break front during the 1920s, and have not reappeared. Minke whales *B. bonaerensis* probably expanded into the blue whale vacated habitat, but were then hunted during the 1970–1980s; their population has since recovered. Some minke whales are now taken in ‘scientific whaling’, twice more from the slope compared to the shelf. Other hunted cetaceans never occurred over the shelf and very few ever occurred in slope waters, and therefore their demise from whaling does not apply to the Ross Sea. No industrial fishing occurred in the Ross Sea until the 1996–1997 summer, when a fishery for Antarctic toothfish *Dissostichus mawsoni* was initiated, especially along the slope. This fishery has grown since then with effects on the ecosystem recently becoming evident. There is probably no other ocean area where the details of biological exploitation can be so elucidated. It appears that the Ross Sea continental shelf remains the least affected of any on the globe. However the same cannot be said of the slope.

Contents

Introduction	233
Definition of the Ross Sea	233
Exploitation	234
Final thoughts	240
Acknowledgements	241
References	241

Introduction

In a study that divided the world ocean into 232 parts and assessed the importance of 17 different anthropogenic factors in altering ecosystem function in each, Halpern and others (2008) indicated that the Ross Sea has been the least impacted of any open ocean, marine area on Earth. Tallying cumulative effects, the Ross Sea received a score of 0.1 compared to the highest scores, which ranged 17.0–19.5 (North Sea, certain equatorial reefs, etc). The impacting factors included widespread pollution of different sorts, artisanal fishing, alien species invasion, climate change (warming, acidification, UV), and others. In regard to commercial fishing, five types were included, but just for the five years from 1999 to 2003. Recent whaling does not seem to have been included.

The case has been made that the Ross Sea is the least affected stretch of ocean remaining on Earth, and therefore deserves protection (Ainley 2002, 2004). Since Halpern and others (2008) did not consider historical data for any region in their analysis (Blight and Ainley 2008), to substantiate that claim about the Ross Sea, a review of the history of earlier exploitation is necessary. Herein, historical and current sealing, whaling, and fishing within the Ross Sea are reviewed, and the case is made that, indeed, the Ross Sea’s continental shelf foodweb

has been until very recently, with the advent of a toothfish fishery, in a state much as it has been over past millennia.

Definition of the Ross Sea

The Ross Sea, about the size of southern Europe, is defined, following the boundary identified by Davey (2004), as the waters overlying the continental shelf and slope extending in a wavering line, including the northward projecting Pennell Bank, from Cape Adare, Victoria Land (71° 17’ S, 170° 14’ E), to Cape Colbeck, Marie Byrd Land (77° 07’ S, 157° 54’ W; Fig. 1). Not included are waters around the Balleny Islands (66° 55’ S, 163° 20’ E), which are the summits of deep-rising sea mounts, 200 km to the northwest of Cape Adare. However some authors have included this area, as well as others north of the Ross Sea shelf and slope (Anon. 2000; Waterhouse 2001; see below).

The Ross Sea is shaped like a right angled triangle, with the height being the north-south oriented Victoria Land coast and the base being the east-west Ross Ice Shelf (or Barrier), extending north and east, respectively from Ross Island (77° 30’ S, 168° 00’ E). To be accurate, the Ross Ice Shelf covers a portion of the Ross Sea basin to the south equal in size to the open-water portion in the north. The mean depth of the exposed shelf is about 500 m, although this varies widely between deep troughs and shallow banks. The shelf break has been defined by some to occur at the level of the troughs, about 800 m, rather than the outward crest of its banks, with the slope descending to 3000 m (Smith and others 2007); the tops of the banks at the shelf break can be <200 m deep. Overall, the Ross Sea bathymetry slopes upward




Fig. 1. The Ross Sea showing locations mentioned in the text. The shelf break, defined as the 800 m isobath, is shown by a dashed line; banks are shown by narrow, solid lines; sea ice is shown as marbled shading in a distribution typical of early December. Discovery Inlet, an embayment in the Ross Ice Shelf that was similar to Bay of Whales, is no longer a feature, having been the product of iceberg calving in the early years of the 20th century. The Balleny Islands are well to the northwest, beyond the boundaries of this map.

from south to north (the banks) owing to the isostatic depression of the continent from the heavy mass of the polar ice cap.

Ross Sea surface water (RSSW) is distinctive and, until recent glacial melting upstream, was the most saline surface water in the Southern Ocean (Jacobs and others 2002). The high salinity of RSSW is related to sea ice formation, with concomitant salt rejection as the ice forms, for much of the year (Jacobs and Comiso 1989). At the Ross Sea (Antarctic) slope front, circumpolar deep water upwells and constrains RSSW to the south (Ainley and Jacobs 1981; Jacobs 1991). The central portion of the Ross Sea, from ice shelf front northward, due initially to incessant winds and subsequently to warming, becomes increasingly clear of sea ice as the season progresses, from October to January. This is the Ross Sea polynya and post polynya (Jacobs and Giulivi 1998, 1999).

Biologically, the Ross Sea is divided into two components, the continental shelf (neritic) and the continental slope (pelagic). The former has as its principal, mid-trophic 'forage species', ice or crystal krill *Euphausia crystallorophias* and the Antarctic silverfish *Pleuragramma antarcticum* (Smith and others 2007). It, therefore, differs from the Ross Sea continental slope food web, dominated by Antarctic krill *E. superba*, a food web that was damaged in the past by the mass removal of whales (see below).

Exploitation

Sealing

Major, industrial sealing did not affect the Ross Sea. No evidence exists that fur seals (*Arctocephalus* spp.), the main target species of early sealers, ever bred in any numbers, if at all, in the vicinity. No specimens have ever been found along Ross Sea shores, and no reports exist noting their presence at sea in the vicinity. The closest, and virtually only location where Antarctic fur seals *A. gazella* could have bred and still reach the Ross Sea in any numbers would have been the very steep sided Balleny Islands. These emerged sea mounts are surrounded by sea ice for most of the year and, moreover, the Antarctic fur seal is not a 'pack ice associated species' (Siniff and others 2008). No evidence exists that fur seals ever occurred on the Balleny Islands, although they were discovered by John Balleny during a search for new sealing grounds in 1839 (Riffenburgh 2006 I: 123–124), and there were inspections by subsequent early explorers, for example members of James Clark Ross's party who attempted to wade ashore (Ross 1847). No evidence exists to indicate long distance migratory movement in southern hemisphere fur seals, similar for example to that which occurs in the northern fur seal *Callorhinus ursinus*, the females of which winter off Japan and California a few thousand kilometers from their Pribilof breeding site (for example Kenyon and Wilke 1953). Therefore, the

migration of Antarctic fur seals from even the closest breeding colony, Macquarie Island (54°37'S, 158°52'E), lying slightly south of Tasmania and New Zealand, is highly unlikely.

In the case of the southern elephant seal *Mirounga leonina*, as late as the early 1980s after sub-Antarctic breeding populations had recovered from decimation (Laws 1994), low numbers of subadult males hauled out for moult in a number of locations during late summer, for example Ross Island and the Victoria Land coast (Brownell and Ainley 1976; Hall and others 2006), as well on offshore rocks at the Balleny Islands (Riffenburgh 2006 I: 123). No longer do they occur in the Ross Sea as the source population on Macquarie Island seriously decreased once again in the 1980s (Laws 1994), and thus the likely density dependent foraging range of that population appears to have contracted. Assuming that a recent analysis concerning Antarctic southern elephant seal rookeries is correct (Ainley and Blight 2009), the decrease was due to the depletion of their fish prey in sub-Antarctic Macquarie Island's surrounding waters and shelves. Recent tracking of elephant seals from Macquarie Island indicate foraging in waters north of the Ross Sea, as defined above (McConnell and others 1992; Van den Hoff and others 2002; Hindell and others 2003).

The early, 'heroic' expeditions killed, for human and dog food, an unknown number of Weddell seals *Leptonychotes weddellii* in southern McMurdo Sound, but by the 1950s the population was thought to have recovered (Stirling 1971). In the same area, the New Zealand Antarctic Programme then killed 50–100 annually, eventually totaling about 2000 animals, in order to feed sled dogs. From 1957 to 1985; the U.S. programme killed a small (unknown) number for scientific reasons during the first few years of this period (Stirling 1971). The population, in the late 1950s prior to initiation of the New Zealand kill was thought to number close to 3000 individuals. It initially decreased in size, but then began to increase in a density dependent response to the kill, followed by a decrease to near 1500 by the mid-1980s when the killing ended. Relative to the Ross Sea population, this decrease was thought to be local, in McMurdo Sound, although the population is partially sustained by emigration from the greater Ross Sea. Therefore, fluctuation in some sort of density dependent emigration from outside McMurdo Sound was probably involved at times (Testa and others 1987).

Since the end of the seal take, the population of seals in southern McMurdo Sound recovered only slightly, now numbering around 2000. It has not recovered further owing to an ocean regime shift in the mid-1970s and changes of sea ice and/or the foodweb (Ainley and others 2005).

Therefore, at the large scale, the Ross Sea shelf is missing several dozen elephant seals and several hundred Weddell seals from the summer food web. A minimum of 10,000 Weddell seals currently breed (adults) along the Victoria Land coast and outlying islands, with unknown

Table 1. A summary of population estimates of marine mammals in the Ross Sea made on the basis of several cruises during the late 1970s-early 1980s (Ainley 1985).

Weddell seal <i>Leptonychotes weddellii</i>	31,990
Leopard seal <i>Hydrurga leptonyx</i>	7,990
Crabeater seal <i>Lobodon carcinophagus</i>	203,700
Ross seal <i>Ommatophoca rossii</i>	5,050
Elephant seal <i>Mirounga leonina</i>	40
Minke whale <i>Balaenoptera bonaerensis</i>	14,280
Killer whale <i>Orcinus orca</i>	3,440

numbers also in the Bay of Whales and along Cape Colbeck (Siniff and Ainley 2008). In total, including young age classes, as many as 30,000 in total may be present in the Ross Sea (Table 1). In addition, there are a few hundred thousand seals of three other species present in the Ross Sea, and which have never been exploited. Mostly they are associated with the marginal ice zone of the eastern and western Ross Sea and the continental slope (Ainley 1985, 2002; Table 1).

Whaling

Caveats regarding Ross Sea whales

It has been proposed that while the extraction of whales, especially blue whales *Balaenoptera musculus intermedia*, from the continental slope of the Ross Sea was important, little whaling was carried out in waters overlying the shelf itself (Ainley 2002, 2003), where only minke whales *B. bonaerensis*, killer whales *Orcinus orca* and Arnoux's beaked whales *Berardius arnuxii* are known to occur at present (Ainley 1985; Ponganis and Kooyman 1995; Van Dam and Kooyman 2004) and, as will be argued below, constitute the neritic cetacean community of the past. Various persons, many privately, have questioned how whalers could have avoided the Ross Sea neritic system, as doing so seemingly would be contrary to the literature (for example Pinkerton 2007). However, a closer look at the available information (see below) indicates that very few blue, even fewer fin and certainly no humpback whales occurred over the shelf in the past or at present, unlike in adjacent waters, and in fact very few individuals of these species, if any, were ever taken from Ross Sea neritic waters (as opposed to blue whales and a few fin whales in waters of the continental slope). The subsequent industrial whaling for minke whales during the 1970–1980s was largely confined to waters north, east and west of the Ross Sea, but nevertheless the population since has recovered following the cessation of whaling in the 1980s (Branch 2006; see below). Unlike the great whales, the Antarctic minke, owing to a major portion of the population that resides in the pack ice, was never decimated to the point of near extinction, thus challenging chances for recovery (Ainley and others 2007a).

Three factors confuse the role of whaling in the potential alteration of the Ross Sea neritic ecosystem. First, the whalers separated the Southern Ocean into parts, such as Ross Sea, Amundsen Sea etc, with these

designations applying to any Antarctic waters lying in the respective general vicinity (Tønnessen and Johnsen 1982). Therefore, based on the definition of the Ross Sea stated above, any historical reports that nebulously refer to 'the Ross Sea' need to be considered carefully. Until recently, even the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) referred to the waters south of New Zealand as the Ross Sea. However, waters north of the Ross Sea shelf, being referred to by oceanographers as the Ross Gyre (Jacobs and others 2002), are quite different as is explained above.

The second factor confusing the perception of the level of whaling in the Ross Sea pertains to a location called the Bay of Whales. The uninformed assumption is made that if whales were so numerous to have deserved the naming of a semi-permanent enclave of the Ross Ice Shelf after them (Fig. 1), then how could the mass killing of whales not have occurred there? The Bay of Whales was named by Ernest Shackleton (1909), who actually appears to have encountered a large concentration of killer whales, or at least he commented on the conspicuousness of the dorsal fins of those whales he saw (baleen whales have very small, unremarkable dorsal fins): 'All around us were numbers of great whales showing their dorsal fins as they occasionally sounded . . . We named this place the Bay of Whales, for it was a veritable playground for these monsters' (Shackleton 1909: 50).

In fact, the Ross Sea is the major area where the diminutive type-C (fish eating) killer whale is found (Pitman and Ensor 2003, Pitman and others 2007), and characteristically they occur in large groups sometimes 50–100 or more, especially where fast ice is breaking up (as noted by Ross 1847 1: 244) thus to expose littleharassed fish. Such a density of animals is quite uncharacteristic of foraging baleen whales (Kasamatsu and others 1998). Roald Amundsen (1912), who based his South Pole expedition at the Bay of Whales, concurred with Shackleton, but also seems likely to have been referring to these killer whales, at least judging from their numbers: 'The name of the Bay of Whales . . . is appropriate enough; for from the time of the break-up of the sea-ice this huge inlet in the Barrier forms a favorite playground for whales, of which we often saw schools of as many as fifty disporting themselves for hours together' (Amundsen 1912: 273). This is the behaviour and group size of type-C killer whales (Pitman and Ensor 2003; Pitman and others 2007; Ainley, personal observation).

Ross (1847) had visited what became known as the Bay of Whales in 1841 and 1842 and, although he wrote extensively of the whales seen during his two voyages to the Ross Sea, especially where large numbers were encountered, he made no mention of whales there or within the several days he explored the barrier front. His cruises were late in the season, well after the ice had broken out of the bay. Other subsequent visitors to the Bay of Whales, including three in the last 20 years (Ainley 1985; Ainley and others 2007a), as well as that of the factory ship *James Clark Ross*, which tied up there for

several weeks during the 1923 and 1924 whaling seasons (Barr and Watt 2005), encountered no whales nor any in the nearby Discovery Inlet (Fig. 1). In fact C.A. Larsen commented specifically on this disappointment (Barr and Watt 2005: 288).

The final item fostering lack of clarity in our understanding of the historical occurrence of whales in the Ross Sea is the assumption that the early explorers were knowledgeable about species identifications (Barr and Watt 2005). In fact, rarely did any of the explorers of the heroic age mention whales by species in their logs, and in a number of cases they appeared to be incorrect. For instance, according to Amundsen (1912): ' . . . we did not receive the impression that there was any very great number of them [whales] out in Ross Sea. The species most commonly seen was the Finner; after that the Blue Whale' (Amundsen 1912: 274).

This is a most curious conclusion given the fact that of the 648 whales taken from the Ross Sea (slope; see below) during the first two years of whaling a decade later (1923–1924), only 29 were fin whales (*B. physalus*) and the vast majority were blue whales. In subsequent years, in the region, blue whales taken outnumbered fins 10 to 1 (Tables 2, 3). In Amundsen's case, and in others, these explorers were probably seeing minke whales, which like fin whales are black and are numerous in the Ross Sea (see below); they referred to them as 'finners'. For instance, Ross: ' . . . a great many whales of small size' (Ross 1847 1: 231); "A great number of whales of two different kinds were seen, the larger kind having an extremely long, erect back-fin [killer whale], whilst that of the smaller species was scarcely discernible" (p.244); "A few whales of the 'finner kind' . . . In the evening 'many whales were seen'" (vol. 2, p.145; see above, blue : fin ratio); " . . . a few whales and some finners were also seen during the day (p.195); " . . . but the small fin-backed whale, as also the piebald kind [blue], were numerous along the pack edge" (p.209).

Historical occurrence and take of whales in the Ross Sea; 'heroic' expeditions.

From the quotations above it is clear that minke whales, along with killer whales, were abundant in the inner Ross Sea during historical times, and that blue whales and minke whales were abundant along the slope. According to D.G. Lillie (1912), biologist on Scott's ship *Terra Nova*: 'It is true that only about three species of whalebone whales were recognised south of the pack, but the number of individuals seen daily around the ship was very great. The two commonest species seen were *Balaenoptera Sibbaldi*, the Blue Whale, and *Balaenoptera rostrata*, the Pike Whale' (Lillie 1912: 481).

The third species is the killer whale, and Lillie (1915: 120) later revised his assessment, saying that 'undoubtedly the commonest cetacean in the Ross Sea' was the killer whale, seen almost every day, with minke whale the second most abundant. In support of this revision, Lillie (1915) logged the location of whale sightings over the shelf and slope, respectively, as follows:

Table 2. Summary of information from Tønnessen and Johnsen (1982; their Table 25 and text) with annotations. All numbers, except as noted, for the last total listed and for any whales taken in Balleny Islands area, are of blue whales (see Table 3).

Season	Whales	Comment
1923–1924	221	Ross Sea, factory ship stationed at Discovery Inlet; whaling along slope (of total 10 fin)
1924–1925	427	Ross Sea, factory ship at Possession Island area, slope; short time at Discovery Inlet; whaling along slope (of total 19 fin)
1925–1926	531	Almost entirely Possession Island area, slope [23]
1926–1927	1117	Possession Island (Ross Sea slope) and Balleny Islands area
1927–1928	2012	Same logistical area as previous season.
1928–1929	1742	Most whales from Balleny Islands area, outside of ice belt
1929–1930	61	All from Ross Sea, that is inside ice belt [23], along slope; any others from Balleny Islands area
Total	6111	Number expressly from Ross Sea slope: 1,240 + unknown numbers taken in 1926–1927 and 1927–1928 (when most were from Balleny Islands area)
1929–1931	5223	NE of Balleny Islands; no operation inside of pack ice, that is in the Ross Sea
1923–1931	18,238	Total for Ross sector including outside Ross Sea proper [23] unlisted above

killer whale, 17 and 0 (but he stated that they were too numerous to log regularly); minke whale, 16 and 8; and blue whale, 5 and 3. The cruises included in his report, both that of *Discovery* (using E. Wilson's notes) and of *Terra Nova*, mostly steamed through shelf waters along the Victoria Land coast (one episode along the ice barrier) and spent relatively little time crossing the slope. This cruise pattern would affect perceptions of relative abundance (see below). Scott (1913) further wrote of killer whales that they were '... less common in the pack but numerous on the coasts' (Scott 1913: 52); and Ponting photographed some apparent type-B killer whales and wrote a description of killer whales pursuing a Weddell

seal and its pup (Savours 1975). Lillie (1915) reported no confirmed sightings of fin, sperm *Physeter macrocephalus* or humpback *Megaptera novaeangliae* whales in waters over the shelf or slope, though they were logged further north.

Lillie (1915) recorded the position of sightings of five blue whales in shelf waters, all spaced regularly, very close to and along the front of the Ross Ice Shelf. These appear to be somewhat anomalous occurrences as, today, top trophic species of any kind are scarce in these waters (Ainley 1985) and there is no Antarctic krill, which is considered a key attribute of this whale species' distribution in the Antarctic (see below). Thus, what these

Table 3. Whales taken during historical times from the Ross Sea sector of the Southern Ocean; information contained in the IWC database, courtesy T.A. Branch; compare Table 2 for specifics of locations within this sector.

Year	Floating factory	Blue	Fin	Sperm	Humpback	Sei	Total
1923	<i>JC Ross</i>	211	10	0	0	0	221
1924	<i>JC Ross</i>	408	19	0	0	0	427
1925	<i>JC Ross</i>	523	8	0	0	0	531
1926	<i>CA Larsen</i>	405	45	0	82	0	532
	<i>NT Nielsen-Alonso</i>	426	27	0	0	0	453
	<i>JC Ross</i>	237	17	0	0	0	254
1927	<i>CA Larsen</i>	785	43	0	9	0	837
	<i>NT Nielsen-Alonso</i>	733	22	0	0	0	755
	<i>JC Ross</i>	564	45	0	7	0	616
1928	<i>CA Larsen</i>	644	23	2	13	0	682
	<i>CA Larsen</i>	111	1	0	1	0	113
	<i>NT Nielsen-Alonso</i>	710	18	0	3	1	732
	<i>JC Ross</i>	530	15	0	0	0	545
1929	<i>CA Larsen</i>	733	192	1	166	0	1092
	<i>Kosmos</i>	995	464	2	356	0	1817
	<i>Kosmos</i>	5	0	0	0	0	5
	<i>NT Nielsen-Alonso</i>	509	128	2	105	0	744
1930	<i>NT Nielsen-Alonso</i>	0	0	0	1	0	1
	<i>JC Ross</i>	342	68	1	24	0	435
	<i>JC Ross</i>	0	12	0	2	0	14
	<i>Southern Princess</i>	459	294	0	121	0	874
Totals		9330	1451	8	890	1	11680

whales were doing is unclear but it is reasonable that they were not feeding. It is possible that these were multiple sightings of the same individual(s).

In regard to the overall distribution of cetaceans, it is very much evident in the logs of Ross, who noted many whales in slope waters but specifically mentioned a major decrease when plying waters to the south over the shelf, except in the bordering pack ice (which he tried to avoid). As noted above, Ross reported only the small, fin-back whale (minke) and the killer whale in southern, shelf waters. This pattern corresponds with that seen in modern times, elaborated upon below, that is whales mainly frequenting the marginal ice zone surrounding the eastern, northern and western border of the Ross Sea polynya (Ainley 1985; Karnovsky and others 2007).

Industrial Whaling.

Further information about historical occurrence patterns of whales can be gleaned from the whaling that soon followed the heroic expeditions. In the first year of Ross Sea whaling, 1923, after finding no whales in the Bay of Whales, the factory ship *James Clark Ross* left anchorage in its protected waters and with its five catcher boats steamed north (Barr and Watts 2005: 289). 'By 8.00 p.m. the small flotilla was clear of this belt of ice [at Bay of Whales] and was again steaming across the ice-free waters of the Ross Sea. Just after they left the ice-edge a number of spouts were seen, but to everyone's disgust they were soon identified as those of Minke whales... But early next morning (28 December) numbers of heavy spouts were sighted on the horizon and all the catchers raced off in pursuit.'

Position data showed these latter whales, blues, to be along the Ross Sea slope. Again and again this pattern is repeated in explorer and whaler logs: large numbers of whales over the slope. The flensers on board *James Clark Ross* complained of working in those slope waters owing to the swell and thus in the end the factory ship often had to remain in Bay of Whales or Discovery Inlet, thereupon having the catchers bring the whales from afar. Larsen noted several times that this slowed operations greatly owing to the long distances that the whales had to be towed (up to 65 km mentioned in one passage) (Barr and Watt 2005).

By the next season, these whalers had deduced the pattern of whale movement, and had also determined the strategy to hunt the whales in the pack ice along the slope (Tønnessen and Johnsen 1982: 351): 'This successful catch was in actual fact the first ice catching on a large scale in the Antarctic, revealing all the necessary criteria: the floating factory was not anchored up or moored to the shore, and catching operations were not carried out in the ice-free Ross Sea, but in the open water between the ice floes.'

Eric Marshall (1930), on the factory ship *C.A. Larsen*, in the final year of the (true) Ross Sea whaling (1929–1930) noted, with map to illustrate, that the blue whales were taken on the northward side of an offshore 'bank'.

In fact, the northern edge of this bank, now known to be several banks running southwest to northeast, constitutes the shelf break (Fig. 1).

The early history of 'Ross Sea' whaling is summarised in Table 2, which gives an idea of the abundance, species composition and distribution of the species that were hunted. In the first year, the effort was entirely contained in the central portion of the Ross Sea slope (all blues, a very few fin whales), with the single factory ship stationed from time to time at the Ross Ice Shelf. In the next two years, almost all of the effort occurred in the vicinity of Possession Island (Fig. 1), at the extreme western end of the Ross Sea shelf break (almost all blues, a few fins). In the next three years, with two then three factory ships added (Table 3), the effort shifted farther west to the waters off Adélie Land (Balleny Islands), but still in the Ross Sea 'sector' (blues, fins, humpbacks and an occasional sei whale *B. borealis*). A final attempt was made in 1929–1930 to enter the Ross Sea for whales, that is the *James Clark Ross* cruise on which Marshall participated (above), but only a few were taken (blues). In the 1950s, factory ships twice attempted to whale in the Ross Sea, south of the ringing ice field, but quickly departed taking few, if any, whales (Tønnessen and Johnsen 1982).

In the 1926–1927 season a pattern was observed that is curious, and may not be 'real', but which colours perceptions of Antarctic whale occurrence even today, and is related to the relationship between ships (killer or survey) and certain whale species in regard to pack ice. In that season the *James Clark Ross* reached the pack ice edge on 12 December, but found the ice belt to be 1000 km wide. This was much wider than in earlier years. It took 21 days to push through but once inside they found few blue whales, no doubt because the ship was then over the shelf. The conclusion was made that the whales, like the ships, had trouble getting through the pack (!), rather than that the whales were over the continental slope, as always. The slope's pack ice cover thus hid the whales from the whalers, who had no capability to search to find them efficiently, particularly because their numbers by then were low anyway. In contrast, in the next season the pack ice belt was unusually narrow, and blue whales were found. In current years, IWC (International Whaling Commission) delegates are being challenged with acknowledging that a great many minke whales occur within the ice pack, where most survey vessels do not venture (Ainley and others 2007a).

Finally, regarding southern right whales *Eubalaena australis* and sperm whales, Ross (1847) is the only person to have ever seen right whales in the southern Ross Sea sector, and in fact well north of the Ross Sea, at latitude 63° 20'S, 174° 30'E, at the north edge of the ice pack. He recorded '... a great many whales, chiefly common black kind, greatly resembling but said to be distinct from the Greenland whale; sperm as well as hunchbacked also observed' (Ross 1847 1: 169).

We can only assume that Ross was familiar with this species (and not so much the others). After all he sailed

extensively in the eastern Canadian Arctic where right whales once abounded, and the reason no one has since seen them in Ross Gyre waters is that they were decimated in their breeding grounds around New Zealand.

Ross also reported sperm whales in a few instances, in each case, as with Lillie (1915), well north of the Ross Sea (waters of the outer slope and deeper; consistent with Nemoto and others 1988). The IWC database indicates that 8 were taken in the region during the 1920s (Table 3). We can assume that their numbers now, in waters north of the Ross Sea, are much reduced as their range contracted upon the population being decimated by whaling farther north (see Whitehead 2000).

Modern occurrence of whales in the Ross Sea

The first modern survey of cetaceans in the Ross Sea was conducted during December and January 1976–1980 (Ainley 1985), much earlier in the season than more recent surveys. The entire Ross Sea was surveyed including pack ice covered portions (3,196 km of track), as the survey vessels were icebreakers that were following a pre-determined track in an oceanographic survey; the ice configuration was similar to that shown in Fig. 1. During this effort, 78 sightings of minke whales, numbering 153 individuals, and 6 sightings of killer whales, numbering 64 whales, were logged. Some unidentified beaked whales were seen on one occasion, very likely Arnoux's beaked whales (Ponganis and Kooyman 1995). From these numbers, it was estimated that about 14,277 minke and 3,442 killer whales were in the Ross Sea (shelf and slope) at that time (Table 1). The killer whale result was thought to be an over estimate due to their propensity to be attracted to the survey vessel (Ainley 1985). In contrast, minke whales were wary of the ship, perhaps because whaling for them was continuing at the time (Ainley and others 2007a).

These same species of whales were sighted on a cruise in May 1998 that constitutes the other seasonal extreme of Ross Sea surveys (Van Dam and Kooyman 2004). All other cetacean surveys of the Ross Sea in recent decades have been conducted in January–February and are all IWC/IDWR–SOWER efforts (see below). The overall pattern in these surveys, similar to that observed by the early explorers, was a preponderance of cetaceans over the slope and within the marginal ice zones of the Ross Sea polynya, with very few over the central shelf (see maps in Ainley 1985; Ichii and others 1998; Karnovsky and others 2007).

In regard to blue whales, on the basis of all available catch, sightings, strandings, *Discovery* marks and recoveries, and acoustic recording data in the Southern Ocean, it has been concluded that 'Blue whales were generally associated with waters deeper than continental shelves. Shallow-water records were typically from regions with narrow continental shelves. . . In the Antarctic, they were most common on deep continental slopes' (Branch and others 2007: 136–137).

Given that the Ross Sea shelf is the broadest in the Southern Ocean, this statement, therefore, is exactly consistent with the story outlined above. These authors' maps show no blue whales taken from or seen in the neritic waters of the Ross Sea and few even on the outer shelf, except for a tiny 'island' of catches just to the east of Ross Island at the western end of the barrier. This must be a record of catch processing and not capture, as in modern times very few whales of any kind are seen in this locality. The waters there have a food web that does not generate high krill or forage fish densities (Ainley 2007; Smith and others 2007) and certainly not *E. superba*, the other attribute of habitat identified by Branch and others (2007) as being critical to blue whale occurrence in the Southern Ocean. In any case, these cases are a few blue whales (see also Lillie observations, above). Finally, Branch and others (2007) noted that blue whales occur in conjunction with oceanographic fronts, and in the Ross Sea, the slope front constitutes an area frequented by Antarctic krill and vast bird numbers (Karnovsky and others 2007). It is a very rich area.

Similarly, a summary of all sighting and capture data for blue, fin and humpback whales for IWC areas III–VI (40°E–150°W), for 1987–2004, reported only three blue whales over the northwestern Ross Sea shelf (Matsuoka and others 2008). The capture data reported was also consistent with the story above: these whales do not and did not occur over the Ross Sea shelf in any numbers (blue), if at all (fin, humpback). In the case of the last species, the pattern is consistent with that described by others (Branch 2009): none in the Ross Sea, shelf or slope, but large numbers off the Adélie Land coast.

Finally, IWC surveys for minke whales (made in open water) indicate them to be abundant in the eastern and western Ross Sea, along the edge of the pack ice, as well as along the slope; few have been found in the south-central Ross Sea (along the Ross barrier) (Ichii and others 1998; Konishi and others 2008). On the basis of existing documentation from ice breaker type survey vessels (Ainley 1985; Ainley and others 2007a), many more would be within the pack ice of the polynya's marginal ice zone.

Since the end of industrial whaling, the minke whale, having not been decimated plus the fact that a significant residual portion of the population frequented the pack ice where whaling ships had difficulty operating (see above) appears to have recovered (Branch 2006). At least, that is, their numbers in the Southern Ocean had reached an asymptote, as indicated by IDCR–SOWER survey results. During the first (1985/86–1990/91) and second (1978/79–1983/84) circumpolar surveys, respectively, 786,000 and 645,000 whales were estimated, which are similar estimates when one takes into consideration the proportion of the Southern Ocean surveyed during the respective surveys. During the most recent survey, however (1991/92–2003/2004), the estimated minke whale numbers (at least in open waters) became much lower (338,000), a state of affairs that has brought much discussion.

Based on IWC abundance estimates by small geographic areas in the most recent circumpolar survey (1991–2003), a population of 21,000 minke whales is estimated for the Ross Sea (T.A. Branch, personal communication, 8 March 2007), including the shelf and slope (but of course any within the pack ice would not have been surveyed). Therefore, this represents an appreciable population increase from the days of minke whaling of the 1970s–early 1980s (see Table 1). It is highly likely that minke whales increased their numbers and expanded into the slope habitat when blue whales were removed in the 1920s. Given density dependent factors they would not be expected to have increased over the shelf, around the periphery of which they already resided. If and when blue whales recover, we might expect them again to frequent slope waters and for minke whales to decrease there. Indeed, in general, minke whales have significantly decreased as the great whales have increased; specifically, this is the most apparent in IWC areas IV and V, the two areas that have shown a dramatic increase in humpback whales (compare Branch 2006, 2009). Humpback and minke whales have been shown to be strong competitors (Friedlaender and others 2008). Blue whales are far too few for analogous studies to be undertaken at present.

Modern Whaling

During the 1970s and 1980s, until the cessation of industrial whaling in 1986, whalers took approximately 19,500 minke whales from IWC areas V and VI; the waters off Adélie Land and Ross Sea ‘sector’ (Brown and Brownell 2001). It is thought that upon the demise of its larger competitors, the Southern Ocean minke whale population had increased overall. For a discussion of possible response of minke whales to competitive release elsewhere, see Laws (1977) and Ballance and others (2006). The minke whale’s population size before whaling is unknown.

Upon the establishment of the whaling moratorium, in 1987, Japan began its ‘scientific whaling’ in waters off east Antarctica, that is IWC areas IV, V and VI. The programme sought to take a total of 440 whales per season from those areas. In portions of areas V-east and VI-west, from a current population of about 21,000 whales (see above), Japan took approximately 440 minke whales from the Ross Sea slope and 295 from the Ross Sea shelf (estimate from Konishi and others 2008: Fig. 1). In the last two seasons, 2006–2008, Japan increased its intended take of minke whales in areas IV, V, and VI to 1,000 whales from the 440 per season taken in 1987–2006. The number taken in the Ross Sea is not publicly known.

Industrial fishing

The world’s continental shelves have largely been depleted of fish, especially demersal species (Hilborn and others 2003; Frank and others 2007). This is true even in the Antarctic. Recent documentation has shown a strong correlation between decreasing populations of fish eating penguins and pinnipeds and the over-exploitation and depletion of fish stocks on the insular shelves and

slopes of low latitude Antarctic islands, including the Antarctic Peninsula (Blight and Ainley 2008; Ainley and Blight 2009). Among the data amassed from the Food and Agriculture Organization (FAO) and CCAMLR, none indicate that industrial fishing had ever occurred in the Ross Sea (Koch 1992), until the last 12 years (Pinkerton and others 2007).

In a process reminiscent of ‘fishing down the food web’ (Pauly and others 1998), led by New Zealand, CCAMLR initiated a programme under which the Antarctic toothfish *Dissostichus mawsoni* biomass will be reduced by 50% from pre-exploitation levels within 35 years, beginning in 1996 (Pinkerton and others 2007). In 1996, an ‘experimental’ fishery began when New Zealand vessels extracted 1 mt, growing to 745 mt by 1999 (data from CCAMLR 2008). Some portion of this catch was from seamounts north of the Ross Sea. In 2000 no longer an experimental fishery, vessels from other countries joined, along with illegal vessels. Once CCAMLR raised the allowable catch from ~2500 mt to ~3500 mt in 2002, vessels from more than 10 countries were attracted and the catch began to meet the quota. CCAMLR believes that as of 2007, and based on the legal take, a 15% reduction in pre-fished biomass has already been achieved (Pinkerton and others 2007).

Although the catch per unit effort in the fishery has not shown any signs of depletion, it is no longer possible to catch adult fish in the southernmost reach of the Ross Sea shelf, in McMurdo Sound, judging from a record that spans 35 years, 1971–2006. Decreases are already evident in the prevalence of toothfish eating killer whales (DeVries and others 2008). In the DeVries and others study more than 4,000 fish were tagged and released, with a few dozen taken over the years for physiological studies.

Final thoughts

The foodweb of the Ross Sea continental shelf, unlike that of the slope, appears to have been largely untouched directly by humans up until the past 10 years. There are fewer elephant seals, formerly rare in the Ross Sea, and fewer Weddell seals in McMurdo Sound (~5% of the shelf area). The cetacean fauna appears to be much as it has been since James Clark Ross first sailed into these waters in the 19th century. It is highly likely that minke whales increased their numbers and expanded into the slope habitat when blue whales were removed in the 1920s, but given density dependent factors they would not be expected to have increased over the shelf where they were already resident. If and when blue whales recover, we might expect them again to frequent slope waters and for minke whales to decrease there.

One major caveat to the apparent near pristine nature of the Ross Sea shelf ecosystem exists. It is possible that the large take of minke whales, especially in pelagic waters north and west of the Ross Sea (IWC area V, VI), during the 1970s – early 1980s (see above), may have

facilitated, along with climate driven changes in Ross Sea polynyas, the increase of Adélie penguins *Pygoscelis adeliae* in the Ross Sea at that time (Ainley and others 2007b). With the cessation of industrial whaling, the penguin increase leveled off. Recent work has shown that the penguins and these whales are direct trophic competitors (Ainley and others 2006), and that Ross Sea penguins spend their non breeding season in the area from which most of the whales were removed, that is north of the Ross Sea near the large scale pack ice edge (Ballard and others in review). This would seem to be, therefore, the major anthropogenic influence so far in altering the Ross Sea neritic foodweb: more penguins in response, partially, to fewer whales in waters of the penguins' non breeding grounds.

It should be noted that this was a case of intense industrial extraction occurring at a time of rapidly changing climate (Ainley and others 2005) and, therefore, it is no longer possible to determine the relative importance of either factor to explain the penguin trends. This is the state of affairs currently for most of the Southern Ocean, in which many populations are decreasing or recovering from heavy exploitation (see reviews in Ballance and others 2006; Koch 1992; Ainley and Blight 2009). Without protection from further industrial fishing and whaling, the Ross Sea neritic foodweb is expected to exhibit the sorts of trophic cascades that have been seen elsewhere as stocks of top trophic, long lived, slow to mature, and slow to reproduce minke whales and Antarctic toothfish are depleted (see review in Baum and Worm 2009). If these cascades are allowed to happen, the Ross Sea's trophic structure would no longer provide an example of how Earth's cold-water, neritic food webs once functioned nor would it remain a laboratory in which the effects of climate change could be investigated without other factors confusing or masking climate effects.

Acknowledgements

This paper was prepared under NSF grant OPP 0440643, but the thoughts expressed are not necessarily those of the National Science Foundation. The paper was greatly improved by the comments of T.A. Branch, R.L. Brownell and P. Penhale. The first and second persons named also provided access to International Whaling Commission databases. I thank them for their help.

References

- Ainley, D.G. 1985. The biomass of birds and mammals in the Ross Sea, Antarctica. In: Siegfried, W.R., P.R. Condy, and R.M. Laws (editors). *Antarctic nutrient cycles and food webs*. Berlin: Springer-Verlag: 498–515.
- Ainley, D.G. 2002. *The Ross Sea, Antarctica: where all ecosystem processes still remain for study*. Hobart, Tasmania: Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR document WGEMM 02/60).
- Ainley, D.G. 2004. Acquiring a 'base datum of normality' for a marine ecosystem: the Ross Sea, Antarctica. Hobart, Tasmania: Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR document WG-EMM-04/20).
- Ainley, D.G. 2007. Insights from the study of the last intact neritic marine ecosystem. *Trends in Ecology and Evolution* 22: 444–445.
- Ainley, D.G., G. Ballard, S. Ackley, L.K. Blight, J.T. Eastman, S.D. Emslie, A. Lescroël, S. Olmastroni, S.E. Townsend, C.T. Tynan, P. Wilson, and E. Woehler. 2007b. Paradigm lost, or, is top-down forcing no longer significant in the Antarctic marine ecosystem? *Antarctic Science* 19: 283–290.
- Anon. (Anonymous). 2000. *Management plan for specially protected area (SPA 4) Balleny Islands, northern Ross Sea, Antarctica*. Wellington, New Zealand: Ministry of Fisheries.
- Ainley, D.G., G. Ballard, and K.M. Dugger. 2006. Competition among penguins and cetaceans reveals trophic cascades in the Ross Sea, Antarctica. *Ecology* 87: 2080–2093.
- Ainley, D.G., and L.K. Blight. 2009. Ecological repercussions of historical fish extraction from the Southern Ocean. *Fish and Fisheries* 10: 13–38.
- Ainley, D.G., E.D. Clarke, K. Arrigo, W.R. Fraser, A. Kato, K.J. Barton, and P.R. Wilson. 2005. Decadal-scale changes in the climate and biota of the Pacific sector of the Southern Ocean, 1950s to the 1990s. *Antarctic Science* 17: 171–182.
- Ainley, D.G., K.M. Dugger, V. Toniolo, and I. Gaffney. 2007a. Cetacean occurrence patterns in the Amundsen and southern Bellingshausen Sea sector, Southern Ocean. *Marine Mammal Science* 23: 287–305.
- Amundsen, R. 1912. *The South Pole, an account of the Norwegian Antarctic expedition in the 'Fram', 1910–1912*. London: John Murray (reprint New York University Press 2001).
- Ballance, L., R.L. Pitman, R.P. Hewitt, D.B. Siniff, W.Z. Trivelpiece, P.J. Clapham, and R.L. Brownell, Jr. 2006. The removal of large whales from the Southern Ocean: evidence for long-term ecosystem effects? In: Estes, J.A., D.P. Demaster, D.F. Doak, T.E. Williams, and R.L. Brownell, Jr. (editors). *Whales, whaling and ocean ecosystems*. Berkeley: University of California Press: 215–230.
- Ballard, G., V. Toniolo, D.G. Ainley, C.L. Parkinson, P. Trathan, and K. Arrigo. in review. Responding to climate change: Adélie penguins confront astronomical and ocean boundaries. *Ecology*.
- Barr, W., and J.P.C. Watt. 2005. Pioneer whalers in the Ross Sea, 1923–33. *Polar Record* 41(219): 281–304.
- Baum, J.K., and B. Worm. 2009. Recent advances in understanding cascading effects of changing oceanic predator abundances. *Journal of Animal Ecology* doi: 10.1111/j.1365-2656.2009.01531.x
- Blight, L.K., and D.G. Ainley. 2008. Southern Ocean not so pristine. *Science* 321: 1443.
- Branch, T.A. 2006. *Abundance estimates for Antarctic minke whales from three completed circumpolar sets of surveys, 1978/79 to 2003/04*. Cambridge, UK: International Whaling Commission (paper SC/58/IA18).
- Branch, T.A. 2009 (in press). Humpback abundance south of 60S from three completed sets of IDCR/SOWER circumpolar surveys. *Journal of Cetacean Research and Management*.

- Branch, T.A., K.M. Stafford, D.M. Palacios, C. Allison, J.L. Bannister, C.L.K. Burton, E. Cabrera, C.A. Carlson, B. Galletti Vernazzani, P.C. Gill, R. Hucke-Gaete, K.C.S. Jenner, M.-N. M. Jenner, K. Matsuoka, Y.A. Mikhalev, T. Miyashita, M.G. Morrice, S. Nishiwaki, V.J. Sturrock, D. Tormosov, R.C. Anderson, A.N. Baker, P.B. Best, P. Borsa, R.L. Brownell, Jr, S. Childerhouse, K.P. Findlay, T. Gerrodette, A.D. Ilangakoon, M. Joergensen, B. Kahn, D.K. Ljungblad, B. Maughan, R.D. Mccauley, S. Mckay, T.F. Norris, Oman Whale and Dolphin Research Group, S. Rankin, F. Samaran, D. Thiele, K. Van Waerebeek, and R.M. Warneke. 2007. Past and present distribution, densities and movements of blue whales *Balaenoptera musculus* in the Southern Hemisphere and northern Indian Ocean. *Mammal Review* 37: 116–175.
- Brown, M.R., and R.L. Brownell. 2001. *Review of catches of great whales taken in the proposed South Pacific sanctuary region*. Cambridge, UK: International Whaling Commission (paper SC52/033).
- Brownell, R.L., and D.G. Ainley. 1976. Southern elephant seals in the Ross Sea. *Antarctic Journal of the United States* 11: 189–190;
- CCAMLR (Commission for the Conservation of Antarctic Marine Living Resources). 2008. Report of the CCAMLR Performance review panel. Hobart, Tasmania: Commission for the Conservation of Antarctic Marine Living Resources.
- Davey, F.J. 2004. Ross Sea bathymetry, 1:2000000, version 1.0. Institute of Geological and Nuclear Sciences, Geophysical Map 16. Lower Hutt, New Zealand: Institute of Geological and Nuclear Sciences.
- DeVries, A.L., D.G. Ainley, and G. Ballard. 2008. *Decline of the Antarctic toothfish and its predators in McMurdo Sound and the southern Ross Sea, and recommendations for restoration*. Hobart, Tasmania: Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR Document EMM 08/21).
- Frank, K.T., B. Petrie, and N.L. Shackell. 2007. The ups and downs of trophic control in continental shelf ecosystems. *Trends in Ecology and Evolution* 22: 236–242.
- Friedlaender, A.S., G.L. Lawson, and P.N. Halpin. 2008. Evidence of resource partitioning between humpback and minke whales around the western Antarctic Peninsula. *Marine Mammal Science* doi: 10.1111/j.1748-7692.2008.00263.x
- Halpern, B.S., S. Walbridge, K.A. Selkoe, C.B. Kappel, F. Micheli, C. D'Agrosa, J.F. Bruno, K.S. Casey, C. Ebert, H.E. Fox, R. Fujita, D. Heinemann, H.S. Lenihan, E.M.P. Madin, M.T. Perry, E.R. Selig, M. Spalding, R. Steneck, and R. Watson. 2008. A global map of human impact on marine ecosystems. *Science* 319: 948–951.
- Hall, B.L., A.R. Hoelzel, C. Baroni, G.H. Denton, B.J. Le Boeuf, B. Overturf, and A.L. Töpf. 2006. Holocene elephant seal distribution implies warmer-than-present climate in the Ross Sea. *Proceedings of the National Academy of Sciences* 103: 10213–10217.
- Hilborn, R., T.A. Branch, B. Ernst, A. Magnusson A., C.V. Minto-Vera, M.D. Scheuerell, and J.L. Valero. 2003. State of the world's fisheries. *Annual Review of Environmental Research* 28:359–99.
- Hindell, M.A., C.J.A. Bradshaw, M.D. Sumner, K.J. Michael, and H.R. Burton. 2003. Dispersal of female southern elephant seals and their prey consumption during the austral summer: relevance to management and oceanographic zones. *Journal of Applied Ecology* 40: 703–715.
- Ichii, T., N. Shinohara, Y. Kujise, S. Nishiwaki, and K. Matsuoka. 1998. Interannual changes in body fat condition index of minke whales in the Antarctic. *Marine Ecology Progress Series* 175: 1–12.
- Jacobs, S.S. 1991. On the nature and significance of the Antarctic slope front. *Marine Chemistry* 35: 9–24.
- Jacobs, S.S., and J. Comiso. 1989. Sea ice and oceanic processes on the Ross Sea continental shelf. *Journal of Geophysical Research* 94 (C12): 18195–18211.
- Jacobs, S.S., and C.G. Giulivi. 1998. Interannual ocean and sea ice variability in the Ross Sea. *Antarctic Research Series* 75: 135–150;
- Jacobs, S.S., and C.G. Giulivi. 1999. Thermohaline data and ocean circulation on the Ross Sea continental shelf. In: Spezie, G., and G. Manzella (editors). *Oceanography of the Ross Sea – Antarctica*. Berlin: Springer-Verlag: 3–16.
- Jacobs, S.S., C.G. Giulivi, and P.A. Mele. 2002. Freshening of the Ross Sea during the late 20th century. *Science* 297: 386–389.
- Kasamatsu, F., P. Ensor, and G.G. Joyce. 1998. Clustering and aggregations of minke whales in the Antarctic feeding grounds. *Marine Ecology Progress Series* 168: 1–11.
- Karnovsky, N., D.G. Ainley, and P. Lee. 2007. The impact and importance of production in polynyas to top-trophic predators: three case histories. In: Smith, W.O., Jr., and D.G. Barber (editors). *Polynyas: windows to the world*. London: Elsevier Publishers: 391–410.
- Kenyon, K.W., and F. Wilke. 1953. Migration of the northern fur seal, *Callorhinus ursinus*. *Journal of Mammalogy* 34: 86–98
- Kock, K.H. 1992. *Antarctic fish and fisheries*. Cambridge and New York: Cambridge University Press.
- Konishi, K., T. Tamura, R. Zenitani, T. Bando, H. Kato, and L. Walløe. 2008. Decline in energy storage in the Antarctic minke whale (*Balaenoptera bonaerensis*) in the Southern Ocean. *Polar Biology* doi 10.1007/s00300-008-0491-3.
- Laws, R.M. 1977. The significance of vertebrates in the Antarctic marine ecosystem. In: Llano, G.A. (editor). *Adaptations within Antarctic ecosystems*. Houston: Gulf Publishing Company: 411–438.
- Laws, R.M. 1994. History and present status of southern elephant seal populations. In: LeBoeuf, B.J., and R.M. Laws (editors). *Elephant seals: population ecology, behaviour and physiology*. Berkeley: University of California Press: 49–65.
- Lillie, D.G. 1913. Summary of biological work carried out on board the *Terra Nova*, 1910–1913. In: Scott, R.F., *Scott's last expedition*. London: Smith, Elder and Co: 475–484.
- Lillie, D.G. 1915. Cetacea. British Antarctic (*Terra Nova*) Expedition, 1910. London: British Museum (Natural history report, zoology 1 (3)): 85–124.
- Marshall, E.H. 1930. Report on a visit to the Ross Dependency. *Geographical Journal* 75: 244–251.
- McConnell, B.J., C. Chambers, and M.A. Fedak. 1992. Foraging ecology of southern elephant seals in relation to the bathymetry and productivity of the Southern Ocean. *Antarctic Science* 4: 393–398.

- Matsuoka, K., T. Hakamada, H. Kiwada, H. Murase, and S. Nishiwaki. 2008. *Distributions and standardized abundance estimates for humpback, fin and blue whales in the Antarctic Areas III E, IV, V and VI W (35° E -145° W), south of 60° S*. Cambridge, UK: International Whaling Commission (paper SC/D06/J7).
- Nemoto, T., M. Okiyama, N. Iwasaki, and T. Kikuchi. 1988. Squids as predators on krill (*Euphausia superba*) and prey for sperm whales in the Southern Ocean. In: Sahrhage, K. (editor). *Antarctic Ocean and Resources Availability*. Berlin: Springer-Verlag: 291–296.
- Pauly, D., V. Christiansen, J. Dalsgaard, R. Froeser, and F. Torres, Jr. 1998. Fishing down marine food webs. *Science* 279: 860–863.
- Pinkerton, M., S. Hanchet, and J. Bradford-Grieve. 2007. Finding the role of Antarctic toothfish in the Ross Sea ecosystem. *Water and Atmosphere* 15: 20–21.
- Pitman, R.L., and P. Ensor. 2003. Three forms of killer whales (*Orcinus orca*) in Antarctic waters. *Journal of Cetacean Research and Management* 5: 1–9.
- Pitman, R.L., W.L. Perryman, D. Leroi, and E. Eilers. 2007. A dwarf form of killer whale in Antarctica. *Journal of Mammalogy* 88: 43–48.
- Ponganis, P.J., and G.L. Kooyman. 1995. Multiple sightings of Arnoux's beaked whales along the Victoria Land coast. *Marine Mammal Science* 11: 247–250.
- Riffenburgh, B. (editor). 2006. *Encyclopedia of the Antarctic*. Abingdon, UK: Routledge.
- Ross, J.C. 1847. *A voyage of discovery and research in the southern and Antarctic regions during the years 1839–1841*. London: John Murray (reprint London: David and Charles reprints 1969).
- Savours, A. 1975. *Scott's last voyage through the Antarctic camera of Herbert Ponting*. New York: Praeger Publishers.
- Scott, R.F. 1913. *Scott's last expedition*. London: Smith, Elder and Co.
- Shackleton, E. 1909. *The heart of the Antarctic, the farthest south expedition, 1907–1909*. London: William Heineman (reprint New York: Penguin Putnam 2000).
- Siniff, D., and D.G. Ainley. 2008. *Aerial surveys of Weddell seals during 2007–08, with notes on the history of aerial censuses in the Ross Sea and recommendations for continued count effort*. Hobart, Tasmania: Commission for the Conservation of Antarctic Marine Resources (CCAMLR Document EMM 08/23).
- Siniff, D.B., J.R.A. Garrott, J.J. Rotella, W.R. Fraser, and D.G. Ainley. 2008. Projecting the effects of environmental change on Antarctic seals. *Antarctic Science* 20: 425–435.
- Smith, W.O., Jr., D.G. Ainley, and R. Cattaneo-Vietti. 2007. Marine ecosystems: the Ross Sea. *Philosophical Transactions of the Royal Society B* 362: 95–111.
- Stirling, I. 1971. Population aspects of Weddell seal harvesting at McMurdo Sound, Antarctica. *Polar Record* 15: 652–667.
- Testa, J.W., and D.B. Siniff. 1987. Population dynamics of Weddell seals (*Leptonychotes weddelli*) in McMurdo Sound, Antarctica. *Ecological Monographs* 57: 149–165.
- Tønnessen, J.N., and A.O. Johnsen. 1982. *The history of modern whaling*. London: C. Hurst and Co.
- Van Dam, R.P., and G.L. Kooyman. 2004. Latitudinal distribution of penguins, seals and whales observed during a late autumn transect through the Ross Sea. *Antarctic Science* 16: 313–318.
- Van Den Hoff, J., H.R., Burton, M.A., Hindell, M.D. Sumner, and C.R. McMahon. 2002. Migrations and foraging of juvenile elephant seals from Macquarie Island within CCAMLR managed areas. *Antarctic Science* 14: 134–145.
- Waterhouse, E. (editor). 2001. *Ross Sea region: a state of the environment report for the Ross Sea region of Antarctica*. Christchurch: New Zealand Antarctic Institute.
- Whitehead, H. 2000. Density-dependent habitat selection and the modeling of sperm whale (*Physeter macrocephalus*) exploitation. *Canadian Journal of Fisheries and Aquatic Science* 57: 223–230.